

Footprints of sticky motion in the phase space of higher dimensional nonintegrable symplectic systems

C. Manchein¹, M. W. Beims^{1,2} and J. M. Rost²

¹Departamento de Física, Universidade Federal do Paraná, 81531-980 Curitiba, PR, Brazil

²Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Strasse 38, D-01187 Dresden, Germany

E-mail: mbeims@fisica.ufpr.br

Abstract. “Sticky” motion in mixed phase space of conservative systems is difficult to detect and to characterize, in particular for high dimensional phase spaces. Its effect on quasi-regular motion is quantified here with four different measures, related to the distribution of the finite time Lyapunov exponents. We study systematically standard maps from the uncoupled two-dimensional case up to coupled maps of dimension 20. We find that sticky motion in all unstable directions above a threshold K_d of the nonlinearity parameter K for the high dimensional cases $d = 10, 20$. Moreover, as K increases we can clearly identify the transition from quasiregular to totally chaotic motion which occurs simultaneously in all unstable directions. The results show that all four statistical measures sensitively probe sticky motion in high dimensional systems.

PACS numbers: 05.45.Jn, 05.45.Pq, 05.45.Ra

Submitted to: *New J. Phys.*

1. Introduction

The dynamics of conservative systems consists of different types of motion, from regular over quasi-regular to chaotic, depending on the perturbation parameter. Usually, in $d = 2$ dimensional conservative systems regular islands (Kolmogorov-Arnold-Moser KAM tori) can break as the perturbation parameter increases, and chaotic trajectories may coexist with regular islands [1]. This regime is called quasiregular. Chaotic trajectories may be trapped around stable islands creating “sticky” motion (for a review see [2]). In such cases the dynamics in phase space can be very structured and the motion is non-ergodic. Analytical results for physical quantities in the quasiregular regime are rare and it is very desirable to obtain tools which quantify degrees of ergodicity and hyperbolicity, as for example the angle between the stable and unstable manifolds obtained via the covariant Lyapunov vectors [3, 4]. Even for apparently totally chaotic systems, tiny regular islands may appear which induce sticky motion as exemplified in Fig. 1a with the phase space of the $d = 2$ dimensional kicked rotor [Eqs. (1) for the nonlinearity parameter $K = 5.79$. The dynamics looks totally chaotic – yet, tiny regular islands appear near the points $(x, p) \sim (4.6, 2.9), (1.7, -2.9)$ (see boxes), if enough initial conditions are sampled. Such islands become visible with sufficient magnification (Figs. 1b-c). Hence, the question arises how these tiny regular islands can




Figure 1. Phase space for $K = 5.79$. (b) and (c) are magnifications from the boxes shown in (a).

be identified with not more information than provided by Fig. 1a. In higher dimensions, the information equivalent to Fig. 1a must be cast in a different form since Poincaré

Surfaces of Section (PSS) cannot be constructed. In fact, for $d > 2$ dimensional conservative systems, due to Arnold diffusion, the KAM surfaces do not divide the whole phase space into a set of closed volumes. Although the effect of stickiness of a N -dimensional invariant tori has been analyzed for the asymptotic decay of correlations and anomalous diffusion [5], to the best of our knowledge, the behavior of sticky trajectories in higher-dimensional systems has never been quantified and explored systematically.

In the following we present extensive numerical simulations to detect and quantify sticky motion in the phase space of quasiregular and chaotic conservative systems with up to 20 dimensions. To fulfill this task we analyze small changes in the finite time Lyapunov exponents (FTLE) distribution using four quantities: the variance σ , the skewness κ_3 , the kurtosis κ_4 and P_Λ , the normalized number of occurrences of the most probable finite time Lyapunov exponent. The first three quantities were analyzed in the two-dimensional standard map for the distribution of the finite time stability exponents to detect islands [6] and to study the fluctuations of the mean FTLE for specific trajectories and the corresponding Kolmogorov like entropy [7].

The FTLEs are the average of local divergence of trajectories and for ergodic systems they converge to the LE obtained in the limit $t \rightarrow \infty$, which is independent of initial conditions [8]. In such cases the FTLE distribution has Gaussian form (normal distribution) [9, 10, 11]. However, in quasiregular systems the chaotic trajectories may visit the neighborhood of regular islands, being trapped there for a while, and the FTLEs do not converge quite well. Such small changes in the FTLEs distribution can be used for many purposes: to obtain information about algebraic or exponential stretching in phase space [12, 13], to quantify ergodicity [14, 15], to study stretching processes in chaotic flows [16], to detect the bimodal [17] or multimodal [18] behavior of the local FTLEs, to analyze the time correlations [19, 20, 21] and Poincaré recurrence [21] in coupled maps, to detect the sticky motion [22, 23] and the arise of regular islands [24] for interacting particles in one- and two-dimensional billiards via P_Λ and, as we will see here, to detect and quantify sticky motion in higher-dimensional nonintegrable conservative systems. In addition, FTLEs have been used in dissipative systems to distinguish between strange chaotic and nonchaotic attractors (see [25, 26, 27, 28] for some examples). In these cases the trajectory switches between long times on the original periodic attractor (negative FTLE) and smaller times on the chaotic saddle with positive FTLEs. This is quite the opposite of what is observed in conservative systems, where the trajectory stays longer times along the chaotic trajectory and smaller times near the regular island, which then reduces the FTLE.

Different from the large literature related to the analysis of sticky motion via the distribution of the *local* FTLEs for some specific trajectories, in this work we analyze the FTLEs distribution *after* $n = 10^7$ iteration of the map and over 10^4 initial conditions distributed uniformly in phase space. Our investigations show that even for very large times the FTLEs still depend on initial conditions when sticky motion is present. The paper is organized as follows. In Section 2 we describe the properties of the variance σ and the distribution of the most probable Lyapunov exponent P_Λ for a two-dimensional

conservative system (the standard map) and compare them with higher cummulants (κ_3 and κ_4). In Section 3 we couple N standard maps and by increasing $N = 2, 3, 5, 10$ we analyze carefully the changes observed in the quantities P_Λ , κ_3 and κ_4 . Section 4 summarizes the main results of the paper.

2. The standard map and the FTLE distributions

The standard map is an ideal model for a conservative chaotic system to study the behavior of FTLE distributions, especially if one is interested in an extension to higher dimensions. The original $d = 2$ dimensional map M was proposed by Chirikov [29] and is defined by

$$M : \begin{cases} p_{n+1} = p_n + K \sin(x_n) & \text{mod } 2\pi, \\ x_{n+1} = x_n + p_{n+1} & \text{mod } 2\pi, \end{cases} \quad (1)$$

where p_n and x_n are the dynamical variables of the system after the n th iteration and K is the nonlinear parameter. This map locally approximates more general nonlinear mappings [1]. The FTLE spectrum is determined for a symplectic system using the standard procedure [30, 31], consequently the sum of all FTLEs is zero. This implies that the FTLEs come in pairs $\pm\Lambda_n^{(i)}$ and that the number of zero exponents is even (i is for the i th unstable direction). Hence, it is sufficient to analyze the $\Lambda_n^{(i)} > 0$. The FTLEs can be defined by

$$\Lambda_n^i(\mathbf{x}) = \frac{1}{n} \ln \| M(\mathbf{x}, n) \mathbf{e}_i(\mathbf{x}) \|, \quad (2)$$

with the discrete iterations n . $M(\mathbf{x}, n) = J(\mathbf{x}_{n-1}) \cdots J(\mathbf{x}_1) J(\mathbf{x}_0)$, and $J(\mathbf{x}_n)$ ($n = 0, \dots, n-1$) are the Jacobian matrices determined at each iteration n . $\mathbf{e}_i(\mathbf{x})$ represents the unit vector in the i th unstable direction. These vectors are re-orthonormalized at each iteration and the FTLE spectrum is obtained using Eq. (2). The usual LE is obtained from $\Lambda^i = \lim_{n \rightarrow \infty} \Lambda_n^i(\mathbf{x})$ which is independent of the initial conditions.

Figure 2a shows the FTLEs distributions $P(\Lambda_n, K)$ as a function of the nonlinearity parameter K for 400 initial conditions and up to $n = 10^7$ iterations. For each initial condition we reckon one FTLE Λ_n , which is the time average of all local FTLE along the trajectory. In Fig. 2a the color changes from light to dark (white over yellow and blue to black) with increasing $P(\Lambda_n, K)$. The gray points below the main curve are related to FTLEs from chaotic trajectories which were trapped for a while close to regular islands. For these trapped chaotic trajectories (or “sticky” motion) the *local* FTLE decreases and consequently the final Λ_n also decreases. In such cases the FTLE distribution is not Gaussian anymore but contains interesting information about the dynamics. Figure 2a also reveals that for some specific values of K the amount of sticky trajectories changes, which will be quantified and characterized next. Note, that such information is not provided by the mean FTLE shown in Fig. 2b.




Figure 2. (a) (Color online) Finite-time distribution of the positive Lyapunov exponent $P(\Lambda_n, K)$ calculated over 400 trajectories up to $n = 10^7$ iterations. With increasing $P(\Lambda_n, K)$ the color changes from light to dark (white over yellow and blue to black). (b) The corresponding mean value $\langle \Lambda_n \rangle$.

2.1. Quantitative characterization of the distributions

Clearly, anomalies in the FTLE distribution reflect sticky motion - however, the entire distribution is not a convenient tool to identify and measure sticky motion. Instead, we will analyze the following four measures:

- a.) The number of occurrences of the most probable FTLEs $P(\Lambda_n^p, K) \equiv P_\Lambda(K)$, defined by [22, 24, 23]

$$\left. \frac{\partial P(\Lambda_n, K)}{\partial \Lambda_n} \right|_{\Lambda_n = \Lambda_n^p} = 0, \quad (3)$$

is the value of the maximum of $P(\Lambda_n, K)$ for each K (see Fig. 2). $P_\Lambda(K) = 1$ means that for all initial conditions the FTLEs are equal (within the precision 10^{-3}). Therefore we expect $P_\Lambda(K) = 1$ for a totally chaotic system and large times. For cases of sticky motion appears or if the normal distribution did not converge to the given precision, $P_\Lambda(K) < 1$.

- b.) The variance defined by

$$\tilde{\sigma} = \langle (\Lambda_n - \langle \Lambda_n \rangle)^2 \rangle, \quad (4)$$

which is the second cummulant of the FTLE distribution. It should increase for sticky motion. We will analyze the properties of the relative variance $\sigma = \tilde{\sigma} / \langle \Lambda_n \rangle^2$ which is more appropriate for higher-dimensional systems. In such cases it is possible to detect small differences relative to each unstable direction.

c.) The skewness is defined by

$$\kappa_3 = \frac{\langle (\Lambda_n - \langle \Lambda_n \rangle)^3 \rangle}{\tilde{\sigma}^{3/2}}. \quad (5)$$

It is the third cumulant of the FTLE distribution and detects the asymmetry of the distribution around its mean value. For $\kappa_3 = 0$ we have the regular distribution. Since sticky motion usually reduces the FTLE, the asymmetry of the distribution leads to $\kappa_3 < 0$.

d.) The kurtosis is defined by

$$\kappa_4 = \frac{\langle (\Lambda_n - \langle \Lambda_n \rangle)^4 \rangle}{\tilde{\sigma}^2} - 3, \quad (6)$$

and detects the shape of the distribution. where $\kappa_4 > 0$ indicates that the distribution is flatter than the regular distribution ($\kappa_4 = 0$) while $\kappa_4 < 0$ reveals a sharper distribution than the normal one.

These four quantities will be used to characterize the degree of sticky motion in the transition from low- to higher-dimensional systems. Altogether they should give all relevant informations which can be extracted from the distributions.

2.2. The two-dimensional case ($d = 2$)

To compare with the higher dimensional cases to be discussed later we summarize here $d = 2$ results for the FTLEs distributions of the standard map in a slightly different K regime than considered by [6]. For all simulations we have used 201 values of K in the




Figure 3. (Color online) Deviations from the mean Lyapunov exponent in the interval $K = (7.0, 9.0)$ for the standard map measured with (a) $P_\Lambda(K)$, (b) $\sigma(K)$, (c) $\kappa_3(K)$ and (d) $\kappa_4(K)$, see Eq. (2.1).

interval $K = (7.0, 9.0)$ (stepsize $\Delta K = 0.01$). As can be seen from Fig. 3a P_Λ has many minima in the interval $K = (7.0, 9.0)$. For all minima of P_Λ a certain amount of sticky

motion is expected. This is indeed true, as we have checked constructing the PSS (not shown) for some values of K where minima in P_Λ appear and for all of them we found sticky motion. Comparing P_Λ with Fig. 3b we see that σ has maxima only for values of K in the interval $(7.15, 7.55)$ and close to 8.12, i.e., the variance does not detect sticky motion but gives important additional information, as we will see later. However, skewness (Fig. 3c) and kurtosis (Fig. 3d) detect sticky motion as identified by P_Λ in (Fig. 3a). Both parameters behave as expected. In Fig. 3c we plotted the skewness κ_3 as a function of K . For almost all minima of P_Λ we observe that the skewness (Fig. 3c) is negative. This implies that the sticky motion induces a left asymmetry in the FTLEs distribution. In other words, the tail on the left side of the distribution is larger than the tail on the right side. For almost all minima of P_Λ the kurtosis κ_4 (Fig. 3d) has a maximum. This is expected since a small P_Λ implies widely distributed Λ_n and consequently a relatively flat distribution (large κ_4).

Obviously, in the two-dimensional case P_Λ , and the cummulants κ_3 and κ_4 are sensitive and consistent detectors of sticky phase space trajectories, with the latter two being more sensitive, while P_Λ is less sensitive, but gives additional information (see, e.g., the interval $K = (7.2, 7.5)$), which will become more obvious in the higher dimensional cases.

3. Coupled standard maps

Coupled standard maps are convenient systems to investigate sticky motion in higher dimensional phase space systematically. The equations for N coupled standard maps read

$$S : \begin{cases} p_{n+1}^{(k)} = p_n^{(k)} + K \sin(x_n^{(k+1)} - x_n^{(k)}), \\ x_{n+1}^{(k)} = x_n^{(k)} + p_{n+1}^{(k)}, \end{cases} \quad (7)$$

where $k = 1, \dots, N$. We consider periodic boundary conditions $p^{(N+1)} = p^{(1)}$, $x^{(N+1)} = x^{(1)}$ and unidirectional next nearest neighbor coupling. In Eq. (7) we must use modulus 2π . The model is invariant under a translation of the x -coordinate. For $N = 2$ it is Hamiltonian, symplectic and conserves the total momentum. For $N > 2$ it is just symplectic. The Lyapunov spectrum always possesses two null exponents for $N > 1$.

3.1. $N = 2$ ($d = 4$): The separable case

In this case the system Eq. (7) has one positive, one negative and two zero FTLEs. We show results related to the positive FTLE in Fig. 4. Clearly, the mean value $\langle \Lambda_n \rangle$ increases with K (Fig. 4a). For $N = 2$ the equations of S (7) decouple in center of mass and relative coordinates. The map in the relative coordinate is the standard map but with parameter $2K$. Therefore, the curve in Fig. 4a should match those obtained for the standard map with parameter values $2K = (14, 18)$. To check this we plotted $\langle \Lambda_n \rangle$ in Fig. 4a (dashed line) for the standard map with values $2K = (14, 18)$. Both curves are identical. For all minima of P_Λ (Fig. 4b) the skewness κ_3 (Fig. 4d) and the kurtosis




Figure 4. (Color online) Comparison of (a) $\langle \Lambda_n \rangle$ (the curve $N = 1$ shows results for the standard map calculated at $2K = (14, 18)$, see text) with (b) $P_{\Lambda_n}(K)$, (c) $\sigma(K)$, (d) $\kappa_3(K)$ and (e) $\kappa_4(K)$ in the interval $K = (7.0, 9.0)$ and for $N = 2$.

κ_4 (Fig. 4e) have also minima and maxima, respectively. At the corresponding values of K many trapped trajectories are expected. Since this case is the standard map with larger nonlinearity parameter ($2K$), we do not discuss further details.

3.2. $N = 3$ ($d = 6$): The largest FTLE is most sensitive to sticky motion

The case $N = 3$ has six FTLEs, two positive, two negative and two zero. The analysis will be done for the distributions of the two positive FTLEs which are labeled such that $\Lambda_n^{(1)} > \Lambda_n^{(2)}$. For $N = 3$ the interval $(7.0, 9.0)$ does not exhibit structures in the FTLE distributions pointing to sticky trajectories. Hence, we analyze the interval $K = (0.01, 2.0)$ which is close to the non-chaotic case $K = 0$. Figure 5 shows the mean values of the first and second FTLEs as a function of K . Both FTLEs increase with K , but $\Lambda_n^{(1)}$ increases faster and has a minimum close to $K \sim 0.2$. Close to this value $\Lambda_n^{(2)}$ starts to be finite. Figures 6 and 7 show the quantities (a) $P_{\Lambda}^{(i)}(K)$, (b) $\sigma^{(i)}$, (d) $\kappa_3^{(i)}(K)$ and (d) $\kappa_4^{(i)}(K)$ as a function of K for $\Lambda_n^{(i=1)}$ and $\Lambda_n^{(i=2)}$, respectively. By comparing these plots we see that both distributions detect the sticky motion for $0.1 \lesssim K \lesssim 0.7$. In this region the minimum value of $\kappa_3^{(i)}(K)$ is ~ -50 for $i = 1$ and ~ -30 for $i = 2$. The maximum value of $\kappa_4^{(i)}(K)$ is 4000 for $i = 1$ and ~ 2000 for $i = 2$. For $K \lesssim 0.1$ we observe in Fig. 7 that $P_{\Lambda}^{(2)}(K) = 1$, meaning that all initial conditions leave to the same $\Lambda_n^{(2)} \sim 0.0$. However, $\kappa_3^{(2)}(K)$ and $\kappa_4^{(2)}(K)$ show that the distribution around $\Lambda_n^{(2)} \sim 0.0$ is not Gaussian. By changing $K = 0.01 \rightarrow 0.1$ the distribution changes slightly from




Figure 5. (Color online) Mean values for the two positive FTLEs $\Lambda_n^{(1)} > \Lambda_n^{(2)}$ for $N = 3$.




Figure 6. (Color online) Comparison of (a) $P_A^{(1)}(K)$ with (b) $\sigma^{(1)}(K)$, (c) $\kappa_3^{(1)}(K)$ and (d) $\kappa_4^{(1)}(K)$ in the interval $K = (0.01, 2.0)$ and $N = 3$.

the right asymmetric form $[\kappa_3^{(2)}(0.01) \sim 1]$ to the left asymmetric form $[\kappa_3^{(2)}(0.1) \sim -3]$. The kurtosis is always flat ($\kappa_4^{(2)} > -3$) but approaches a regular distribution ($\kappa = 0$) when $K \sim 0.05$. Similar behavior is observed in Fig. 6 for $\kappa_3^{(1)}(K)$ and $\kappa_4^{(1)}(K)$. For the relative variance is shown to be relevant only at very low values of K (for both FTLEs).

For $K \gtrsim 0.7$ the distribution related to $\Lambda_n^{(1)}$ detects the sticky motion at $K \sim 0.83, 1.40$ and $K \sim 1.32$, while for $\Lambda_n^{(2)}$ it detects $K \sim 1.40$ and $K \sim 0.83$ (but only $\kappa_4^{(2)}(K)$). This shows the intuitively expected result that when the nonlinearity parameter increases the most unstable dimension (related to $\Lambda_n^{(1)}$) is more sensitive to the sticky motion.

3.3. $N = 5$ ($d = 10$): The smallest FTLE defines a critical nonlinearity K_d

In Fig. 8 the mean values for the positive FTLEs $\Lambda_n^{(1)} > \Lambda_n^{(2)} > \Lambda_n^{(3)} > \Lambda_n^{(4)}$ are shown as a function of K . For smaller values of $K \lesssim 0.15$, $\langle \Lambda_n^{(3)} \rangle$ and $\langle \Lambda_n^{(4)} \rangle$ are almost zero, while $\langle \Lambda_n^{(1)} \rangle$ increases initially very fast for $K = 0.01 \rightarrow 0.05$ and then slowly




Figure 7. (Color online) Comparison of (a) $P_{\Lambda}^{(2)}(K)$ with (b) $\sigma^{(2)}(K)$, (c) $\kappa_3^{(2)}(K)$ and (d) $\kappa_4^{(2)}(K)$ in the interval $K = (0.01, 2.0)$ and $N = 3$.




Figure 8. (Color online) Mean values for the four positive FTLEs $\Lambda_n^{(1)} > \Lambda_n^{(2)} > \Lambda_n^{(3)} > \Lambda_n^{(4)}$ for $N = 5$.

for $K = 0.05 \rightarrow 0.15$. For K above 0.2 all mean FTLEs increase linearly with different slopes size-wise ordered as the FTLEs themselves.

Figure 9 shows that for $K \gtrsim 0.3$ the distributions from all FTLEs approach the normal (Gaussian) form expected for a totally chaotic regime in contrast to the cases $N = 1, 2, 3$ discussed before, where sticky motion also occurs for higher values of K . Therefore, there is a clear transition from the quasiregular to the totally chaotic dynamics at $K_c \sim 0.3$ *simultaneously* in all unstable directions. On the other hand, all quantities detect the sticky motion for $K \lesssim 0.25$. Interestingly, above a critical value $K_d \sim 0.13$, the four indicators of sticky motion have a similar qualitative behavior for all FTLEs, i. e., independently of the unstable directions. For example, the distributions for *all* FTLEs have a small left asymmetry ($\kappa_3 \sim -0.75$) with no flatness ($\kappa_4 \sim 0$) at $K = 0.13$, and a left asymmetry ($\kappa_3 \sim -1.50$) with flatness ($\kappa_4 \sim 3.0$) at $K = 0.2$. The critical value $K_d \sim 0.13$ is the point where $P_{\Lambda}^{(4)}(K)$ detects the sticky motion from the distribution of the last (the *smallest*) FTLE [see Fig. 9a]. The value of K_d can also be determined approximately from the onset of linear K -dependence of the mean FTLEs




Figure 9. (Color online) (a) $P_{\Lambda}^{(i)}(K)$, (b) $\sigma^{(i)}(K)$, (c) $\kappa_3^{(i)}(K)$ and (d) $\kappa_4^{(i)}(K)$ in the interval $K = (0.01, 1.0)$, $N = 5$ and for all $i = 1, 2, 3, 4$.

in Fig. 8.

These results show that the sticky motion induces differences between the distributions of $\Lambda_n^{(i)}$ only for $K < K_d$. Since sticky motion is a consequence of regular structures in phase space, this suggests that there could exist disconnected invariant regions in phase-space. In the interval $K_d < K < K_c$ on the other hand, the sticky motion affects all unstable dimensions by the same amount leading to similar distributions for different FTLEs. Finally, in the totally chaotic regime $K > K_c$ all unstable directions are affected proportionally to each other. To explain this better we show in Fig. 10 the time evolution of $\Lambda_n^{(i)}$ ($i = 1, 2, 3, 4$) for one exemplary trajectory in the three distinct regions of K . Figure 10(a) shows the behavior of the FTLEs for $K = 0.6 > K_c$ and no sticky motion is observed since no FTLE goes to zero for some shorter time intervals. For the intermediate case $K_d < K = 0.15 < K_c$ we see in Fig. 10(b) that for times around 0.5×10^6 (see arrow) *all* four FTLEs are zero, so the sticky motion occurs in all unstable directions and a common behavior is observed. Physically what happens is that the linear momentum of *each* site becomes approximately constant during sticky motion. This is clearly shown in Fig.11(a), where

$p_n^{(j)}$ ($j = 1, 2, 3, 4, 5$) is plotted as a function of n . At the sticky time $n \sim 0.5 \times 10^6$ (see black box), $p_n^{(j)}$ is constant for *all* sites. This agrees with the conjecture made




Figure 10. (Color online) Time evolution of $\Lambda_n^{(i)}$ ($i = 1, 2, 3, 4$) for one exemplary trajectory with $N = 5$ and (a) $K = 0.6 > K_c$, (b) $K_d < K = 0.15 < K_c$, (c) $K = 0.05 < K_d$.

by Froeschlé [32] and observed later in coupled maps by [20], i. e., that a conservative system with $N \geq 3$ has N or one (the energy) constants of motion. In other words, following this conjecture when the sticky motion occurs all FTLEs should temporarily be very close to zero, as already observed. However, this changes when the region $K < K_d$ is analyzed, as can be observed in Fig. 10(c) for times around 0.5×10^6 and 0.85×10^6 (see arrows). Only $\Lambda_n^{(1)}$ is larger and the remaining three FTLEs are very close to zero. In this case, at the sticky time $n \sim 0.5 \times 10^6$ (see black box), the momenta $p_n^{(1)}, p_n^{(3)}$ and $p_n^{(5)}$ are not conserved anymore. See Fig. 11(b).

3.4. $N = 10$ ($d = 20$)

The case $N = 10$ has nine positive FTLEs. In this section we analyze again the interval $K = (0.01, 1.0)$ but with 10^3 initial conditions only. In Fig. 12 the mean values for the FTLEs are plotted as a function of K . For $K \gtrsim 0.2$ all mean FTLEs increase linearly and the largest ones increase faster. For $K \lesssim 0.2$ the K dependence is different for different FTLEs. For $K \lesssim 0.1$ almost all (but two $i = 1, 2$) are very close to zero.

Figure 13 compares (a) $P_\Lambda^{(i)}(K)$, (b) $\sigma^{(i)}(K)$, (c) $\kappa_3^{(i)}(K)$ and (d) $\kappa_4^{(i)}(K)$ in the interval $K = (0.01, 1.0)$ for all positive FTLEs ($i = 1 \rightarrow 9$). For $K > K_c \sim 0.28$ all quantities indicate that the distributions from all FTLEs approach the normal form expected for a totally chaotic regime. Therefore, again we have a clear transition from quasi-regular to totally chaotic dynamics at K_c which occurs *simultaneously* in all unstable directions. For $K < K_c$ all quantities detect the sticky motion. Also here




Figure 11. (Color online) Time evolution of $p_n^{(j)}$ ($j = 1, 2, 3, 4, 5$) for the trajectory from fig 10 (a) $K_d < K = 0.15 < K_c$, (b) $K = 0.05 < K_d$. Black boxes show the sticky region.




Figure 12. (Color online) Mean values for the nine positive FTLEs $\Lambda_n^{(1)} > \Lambda_n^{(2)} > \dots > \Lambda_n^{(9)}$, for $N = 10$.

we observe that for K above the critical value $K_d \sim 0.09$ the four quantities behave similarly, independent of the FTLEs. The distributions for all FTLEs have a small left asymmetry ($\kappa_3 \sim -0.75$) with no flatness ($\kappa_4 \sim 0$) at $K = 0.13$, and a left asymmetry ($\kappa_3 \sim -1.50$) with flatness ($\kappa_4 \sim 3.0$) at $K = 0.2$. The critical value $K_d \sim 0.09$ is again exactly the point where $P_\Lambda^{(9)}(K)$ detects the sticky motion from the distribution of the last (the *smallest*) FTLE [see Fig. 9(a)]. It can be observed that the distribution of the largest FTLE is the last (when K increases from 0.01) to “join” the collective behavior for $K_d \gtrsim 0.09$ [please see Fig. 13(b)-(d)]. The value K_d can also be determined approximately from the region in Fig. 12 where the K dependence of the mean FTLEs is not linear.

Figure 14 shows the time evolution of $\Lambda_n^{(i)}$ ($i = 1 \rightarrow 9$) for one exemplary trajectory in the three distinct regions of K . In Fig. 14(a) with $K = 0.6 > K_c$ no sticky




Figure 13. (Color online) Quantities (a) $P_{\Lambda}^{(i)}(K)$, (b) $\sigma^{(i)}(K)$, (c) $\kappa_3^{(i)}(K)$ and (d) $\kappa_4^{(i)}(K)$ as a function of $K = (0.01, 1.0)$, for $i = 1 \rightarrow 9$ and $N = 10$.

motion is observed since no FTLE goes to zero for some shorter time intervals. For the intermediate case $K_d < K = 0.15 < K_c$ we see in Fig. 14(b) the two different scenarios as conjectured. For times around 0.22×10^6 (see arrow) *all* nine FTLEs are close to zero indicating that the sticky motion occurs in all unstable directions. For times around 0.8×10^6 (see arrow) on the other hand, eight FTLEs go to zero but the largest one remains positive. This behavior is observed even more clearly in the region $K < K_d$, see Fig. 14(c). For times in the interval 0.12×10^6 to 0.18×10^6 only $\Lambda_n^{(1)}$ is large and the remaining eight FTLEs are very close to zero.

4. Conclusions

It is well known that the dynamics of conservative systems consists of qualitatively different types of motion, from regular over quasi-regular to chaotic, depending on the perturbation parameter. It is difficult to identify these types of motion in high-dimensional systems, especially in the quasi-regular regime where the sticky motion can suppress Arnold diffusion and distinct stochastic regions with different properties may




Figure 14. (Color online) Time evolution of $\Lambda_n^{(i)}$ ($i = 1 \rightarrow 9$) for one exemplary trajectory with $N = 10$ and (a) $K = 0.6 > K_c$, (b) $K_d < K = 0.15 < K_c$, (c) $K = 0.05 < K_d$.

appear [33]. Therefore, it is very desirable to develop tools which quantify the degree of sticky motion, hyperbolicity and ergodicity. Following ideas from [6] the purpose of the present work is to quantify the degree of sticky motion in the phase space of higher-dimensional systems. To fulfill this task we have systematically analyzed small changes in the FTLE distributions using the skewness κ_3 , which detects the asymmetry of the distribution, the kurtosis κ_4 , which measures the flatness of the distribution and the normalized number of occurrences of the most probable finite time Lyapunov exponent P_Λ , which indicates to which extent all trajectories started are under the influence of the same FTLE.

Using coupled standard maps with nonlinearity parameter K we show that the quantities $\kappa_3(K)$, $\kappa_4(K)$ and $P_\Lambda(K)$ are very sensitive to detect the sticky motion and capable to characterize the dynamics when the dimension d of the system increases ($d = 2, 4, 6, 10, 20$). Besides detecting accurately all cases of sticky motion, we found the following additional interesting results: i) ($d = 6$) the largest FTLE is most sensitive to the sticky motion; ii) ($d = 10, 20$) the smallest FTLE defines the critical nonlinearity parameter K_d , below which disconnected invariant regions in phase-space are expected and $\Lambda_n^{(i)}(K)$ behaves nonlinearly. For $K > K_c$ a clear transition from the quasiregular to the totally chaotic dynamics occurs *simultaneously* in all unstable directions. In the interval $K_d < K < K_c$ all quantities detect the presence of sticky motion and distributions for *different* FTLEs behave *similarly*, meaning that the sticky motion affects all unstable directions proportionally by the same amount and a kind of common behavior of the distributions is observed. Physically what happens is that the linear momentum of *all* sites is constant at the sticky motion.

Close-to-zero FTLEs play also an important role in other higher-dimensional

system, for example in connection to hydrodynamic modes [34, 35] (among others) and the analysis of hyperbolicity [6, 36]. Results of the present work contribute to categorize the rich and complicated quasiregular dynamics of higher-dimensional conservative systems and also to understand how nonlinearity is distributed along different unstable directions [19, 37].

5. Acknowledgments

The authors thank FINEP (under project CTINFRA-1) for financial support. C.M. also thanks CNPq for financial support and Prof. Giulio Casati for the hospitality by the Center for Nonlinear and Complex Systems at the Università dell'Insubria where part of this work was done.

References

- [1] A. J. Lichtenberg and M. A. Lieberman. *Regular and Chaotic Dynamics*. Springer-Verlag, 1992.
- [2] G. M. Zaslavsky. *Hamiltonian chaos and fractional dynamics*. Oxford, University Press, 2008.
- [3] G. Lapeyre. *Chaos*, **12**:688, 2002.
- [4] F. Ginelli, P. Poggi, A. Turchi, H. Chaté, R. Livi, and A. Politi. *Phys. Rev. Lett.*, **99**:130601, 2007.
- [5] E. G. Altmann and H. Kantz. *Europhys. Lett.*, **78**:10008, 2007.
- [6] S. Tomsovic and A. Lakshminarayan. *Phys. Rev. E*, **76**:036207, 2007.
- [7] C. Froeschlé and CH. Froeschlé. *Celestial Mechanics and Dynamical Astronomy*, **56**:307, 1993.
- [8] V. I. Oseledec. *Trans. Moscow Math.*, **19**:197, 1968.
- [9] P. Grassberger, R. Badii, and A. Politi. *J. Stat. Phys.*, **51**:135, 1988.
- [10] E. J. Kostelich, I. Kan, C. Grebogi, E. Ott, and J. A. Yorke. *Physica D*, **109**:81, 1997.
- [11] M. Falcioni, U. Marini Bettolo Marconi, and A. Vulpiani. *Phys. Rev. A*, **44**:2263, 1991.
- [12] M. A. Sepúlveda, R. Badii, and E. Pollak. *Phys. Rev. Lett.*, **63**:1226, 1989.
- [13] M. Ding, T. Bountis, and E. Ott. *Phys. Lett. A*, **151**:395, 1990.
- [14] C. Amitrano and R. S. Berry. *Phys. Rev. Lett.*, **68**:729, 1992.
- [15] C. Amitrano and R. S. Berry. *Phys. Rev. E*, **47**:3158, 1993.
- [16] D. Begie, A. Leonard, and S. Wiggins. *Phys. Rev. Lett.*, **70**:275, 1993.
- [17] J. D. Szezech, S. R. Lopes, and R. L. Viana. *Phys. Lett. A*, **335**:394, 2005.
- [18] M. Harle and U. Feudel. *Chaos, Solitons & Fractals*, **31**:130, 2007.
- [19] P. Grassberger and H. Kantz. *Phys. Lett. A*, **113**:167, 1985.
- [20] H. Kantz and P. Grassberger. *Phys. Lett. A*, **123**:437, 1987.
- [21] R. Artuso and C. Manchein. *Phys. Rev. E*, **80**:036210, 2009.
- [22] M. W. Beims, C. Manchein, and J. M. Rost. *Phys. Rev. E*, **76**:056203, 2007.
- [23] H. A. Oliveira, C. Manchein, and M. W. Beims. *Phys. Rev. E*, **78**:046208, 2008.
- [24] C. Manchein and M. W. Beims. *Chaos Solitons & Fractals*, **39**:2041, 2009.
- [25] X. Wang, M. Zahn, C. Lai, and Y. Lai. *Phys. Rev. Lett.*, **92**:074102, 2004.
- [26] S. S. Negi A. Prasad and R. Ramaswamy. *Int. J. Bifurcation Chaos Appl. Sci. Eng.*, **11**:291, 2001.
- [27] U. Feudel, J. Kurths, and A. S. Pikovsky. *Physica D*, **88**:173, 1995.
- [28] C. Grebogi, E. Ott, S. Pelikan, and J. A. Yorke. *Physica D*, **13**:261, 1984.
- [29] B. V. Chirikov. *Phys. Rep.*, **52**:263, 1979.
- [30] G. Benettin, L. Galgani, A. Giorgilli, and J.-M. Strelcyn. *Meccanica*, **15**(9):09, 1980.
- [31] A. Wolf, J. B. Swift, H. L. Swinney, and J. A. Vastano. *Physica D*, **16**:285, 1985.

- [32] C. Froeschlé. *Phys. Lett. A*, **16**:172, 1972.
- [33] M. Pettine and A. Vulpiani. *Phys. Lett.* , **106A**:207, 1984.
- [34] H. L. Yang and G. Radons. *Phys. Rev. Lett.* , **100**:024101, 2008.
- [35] M. Romero-Bastida and E. Braun. *J. Phys. A*, **41**:375101, 2008.
- [36] J. C. Vallejo, R. L. Viana, and M. A. F. Sanjuan. *Phys. Rev. E*, **78**:066204, 2008.
- [37] C. Manchein, J. Rosa, and M. W. Beims. *Physica D*, **238**:1688, 2009.